


This listing of claims will replace all prior versions, and listings, of claims in the application.

**Listing of Claims:**

1-28. (Canceled)

29. (new) A method for the treatment of a disease or disorder selected from the group consisting of neurodegenerative disease, stroke, and Alzheimer's disease comprising administering to a subject in need of such treatment an effective amount of a compound of the Formula I:



wherein:

W is A-B-D;

A is aryl(CH<sub>2</sub>)<sub>n</sub>, heteroaryl(CH<sub>2</sub>)<sub>n</sub>, alkyl having from one to about 14 carbons, alkenyl having from two to about 14 carbons, or cycloalkyl having from 3 to about 10 carbons, said A group being optionally substituted with one or more J groups;

B is a bond or CO, SO, SO<sub>2</sub>, OCO, NR<sup>5</sup>CO, NR<sup>5</sup>SO<sub>2</sub>, or NR<sup>5</sup>SO;

D is a bond, an amino acid residue, or a peptide composed of 2 to about 5 amino acid residues, said amino acid residue(s) being independently defined by the formula -NH-<sup>\*\*</sup>CH(R<sup>6</sup>)-CO-, in which <sup>\*\*</sup> denotes the α carbon of an α-amino acid residue possessing, when R<sup>6</sup> is other than hydrogen, the D- configuration, the L- configuration, or a mixture of D- and L-;

n is an integer from 0 to about 6;

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are, independently, hydrogen, alkyl having from one to about 14 carbons, or cycloalkyl having from 3 to about 10 carbons, said alkyl, and cycloalkyl groups being optionally substituted with one or more J groups; and

J is halogen, lower alkyl, aryl, heteroaryl, haloaryl, amino optionally substituted with one to three aryl or lower alkyl groups, guanidino, alkoxycarbonyl, amido, lower alkylamido, sulfonamido, lower alkyl sulfonamido, lower alkylsulfonyl, lower alkylsulfoxy, lower alkylthio, lower alkoxy, aryloxy, arylalkyloxy, hydroxy, carboxy, cyano, or nitro; and

\* denotes the  $\alpha$  carbon of an  $\alpha$ -amino acid residue possessing, when  $R^2$  is other than hydrogen, the D- configuration, the L- configuration, or a mixture of the D- and L- configurations.

30. (new) The method of claim 29 wherein  $R^1$  is alkyl or alkyl substituted with J, wherein J is lower alkoxy.

31. (new) The method of claim 30 wherein  $R^1$  is benzyl, methoxymethyl, or butyl.

32. (new) The method of claim 29 wherein  $R^2$  is alkyl or alkyl substituted with J, wherein J is arylalkyloxy or aryl.

33. (new) The method of claim 30 wherein  $R^2$  is isobutyl or benzyloxymethyl.

34. (new) The method of claim 29 wherein  $R^3$  is H.

35. (new) The method of claim 29 wherein  $R^4$  is alkyl, alkyl substituted with J, cycloalkyl, or cycloalkyl substituted with J wherein J is aryl, haloaryl, alkyl or heteroaryl.

36. (new) The method of claim 35 wherein  $R^4$  is methyl, ethyl, propyl, butyl, benzyl, (pentafluorophenyl)methyl, tert-butyl, or 4-methylcyclohexyl.

37. (new) The method of claim 29 wherein W is benzyloxycarbonyl, methanesulfonyl, benzoyl, tert-butoxycarbonyl, or benzyloxycarbonyl-leucyl.

38. (new) The method of claim 29 wherein  $R^3$  is H, and  $R^1$  is alkyl or alkyl substituted with J, wherein J is lower alkoxy.

39. (new) The method of claim 29 wherein  $R^3$  is H, and  $R^2$  is alkyl or alkyl substituted with J wherein J is arylalkyloxy or aryl.

40. (new) The method of claim 29 wherein  $R^3$  is H, and  $R^4$  is alkyl, alkyl substituted with J, cycloalkyl, or cycloalkyl substituted with J wherein J is aryl, alkyl, haloaryl, or heteroaryl.

41. (new) The method of claim 29 wherein  $R^3$  is H,  $R^1$  is alkyl or alkyl substituted with J wherein J is lower alkoxy, and  $R^2$  is alkyl or alkyl substituted with J wherein J is arylalkyloxy or aryl.

42. (new) The method of claim 29 wherein  $R^3$  is H,  $R^1$  is alkyl or alkyl substituted with J, wherein J is lower alkoxy, and  $R^4$  is alkyl, alkyl substituted with J, cycloalkyl, or cycloalkyl substituted with J wherein J is aryl, haloaryl, alkyl or heteroaryl.

43. (new) The method of claim 29 wherein  $R^3$  is H,  $R^1$  is alkyl or alkyl substituted with J wherein J is lower alkoxy,  $R^4$  is alkyl, alkyl substituted with J, cycloalkyl, or cycloalkyl substituted with J wherein J is aryl, haloaryl, alkyl or heteroaryl, and  $R^2$  is alkyl or alkyl substituted with J wherein J is arylalkyloxy or aryl.

44. (new) The method of claim 29 wherein  $R^1$  is benzyl, methoxymethyl, or butyl;  $R^2$  is isobutyl or benzyloxymethyl;  $R^3$  is hydrogen;  $R^4$  is methyl, ethyl, propyl, butyl, benzyl, (pentafluorophenyl)methyl, tert-butyl, or 4-methylcyclohexyl; and W is benzyloxycarbonyl, methanesulfonyl, benzoyl, tert-butoxycarbonyl, or benzyloxycarbonyl-leucyl.

45. (new) The method of claim 29 wherein  $R^1$  is benzyl;  $R^2$  is isobutyl; \* denotes the  $\alpha$  carbon of an  $\alpha$ -amino acid residue possessing the L-configuration;  $R^3$  is hydrogen;  $R^4$  is methyl, ethyl, propyl, butyl, benzyl, (pentafluorophenyl)methyl, tert-butyl, or 4-methylcyclohexyl; and W is benzyloxycarbonyl or benzyloxycarbonyl-leucyl.

46. (new) The method of claim 29 wherein  $R^1$  is benzyl;  $R^2$  is benzyloxymethyl; \* denotes the  $\alpha$  carbon of an  $\alpha$ -amino acid residue possessing the D-configuration;  $R^3$  is hydrogen;  $R^4$  is methyl, ethyl, or benzyl; and W is methanesulfonyl.

47. (new) The method of claim 29 wherein W, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are selected in accordance with the following table:

W	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
BnOCO	Bn	L-CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	H	CH <sub>3</sub>
BnOCO	Bn	L-CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	H	CH <sub>2</sub> CH <sub>3</sub>
BnOCO	Bn	L-CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	H	Bn
BnOCO	Bn	L-CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	H	CH <sub>2</sub> C <sub>6</sub> F <sub>5</sub>
BnOCO	Bn	L-CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	H	tBu
BnOCO	Bn	L-CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	H	(4-methyl-cyclohexyl)
CH <sub>3</sub> SO <sub>2</sub>	CH <sub>2</sub> OCH <sub>3</sub>	D-CH <sub>2</sub> OBn	H	Bn
CH <sub>3</sub> SO <sub>2</sub>	Bn	D-CH <sub>2</sub> OBn	H	Bn
CH <sub>3</sub> SO <sub>2</sub>	Bn	D-CH <sub>2</sub> OBn	H	CH <sub>2</sub> CH <sub>3</sub>
BnOCO	Bn	L-CH(CH <sub>3</sub> ) <sub>2</sub>	H	Bn
BnOCO	(CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>	L-CH(CH <sub>3</sub> ) <sub>2</sub>	H	Bn
Cbz-Leu	Bn	L-CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	H	CH <sub>3</sub>
Cbz-Leu	Bn	L-CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	H	Bn
PhCO	(CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>	L-Bn	H	CH <sub>2</sub> CH <sub>3</sub>
BnOCO	Bn	L-CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	H	(CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>

48. (new) The method of claim 29 wherein

W is benzyloxycarbonyl;

R<sup>1</sup> is benzyl;

R<sup>3</sup> is H;

R<sup>2</sup> is L-CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub> or L-CH(CH<sub>3</sub>)<sub>2</sub>;

and R<sup>4</sup> is selected from the group consisting of -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, benzyl, -CH<sub>2</sub>C<sub>6</sub>F<sub>5</sub>, t-butyl, and 4-methylcyclohexyl.

49. (new) The method of claim 29 wherein the disease or disorder is neurodegenerative disease.

50. (new) The method of claim 29 wherein the disease or disorder is stroke.

**DOCKET NO.:** CEPH-2311

**PATENT**

**Application No.:** Not Yet Assigned

**Preliminary Amendment - First Action Not Yet Received**

51. (new) The method of claim 29 wherein the disease or disorder is Alzheimer's disease.